Analgesia Medical Directive

A Primary Care Paramedic may provide the treatment prescribed in this Medical Directive if authorized.

Indications

Pain

Conditions

acetaminophen				
Age	≥ 12 years			
LOA	Unaltered			
HR	N/A			
RR	N/A			
SBP	N/A			
Other	N/A			

	ibuprofen
Age	≥ 12 years
LOA	Unaltered
HR	N/A
RR	N/A
SBP	N/A
Other	N/A

ketorolac				
Age	≥ 12 years			
LOA	Unaltered			
HR	N/A			
RR	N/A			
SBP	Normotension			
Other	N/A			

Contraindications

acetaminophen

Acetaminophen use within previous 4 hours

Allergy or sensitivity to acetaminophen

Hx of liver disease

Active vomiting

Unable to tolerate oral medication

Suspected ischemic chest pain

ibuprofen

NSAID use within previous 6 hours

Allergy or sensitivity to ASA or NSAIDs

Patient on anticoagulation therapy

Current active bleeding

Hx of peptic ulcer disease or GI bleed

Pregnant

If asthmatic, no prior use of ASA or other NSAIDs

CVA or TBI in the previous 24 hours

Known renal impairment

Active vomiting

Unable to tolerate oral medication

Suspected ischemic chest pain

ketorolac

NSAID use within previous 6 hours

Allergy or sensitivity to ASA or NSAIDs

Patient on anticoagulation therapy

Current active bleeding

Hx of peptic ulcer disease or GI bleed

Pregnant

If asthmatic, no prior use of ASA or other NSAIDs

CVA or TBI in the previous 24 hours

Known renal impairment

Suspected ischemic chest pain

Treatment

Consider acetamino	ohen	
	Age	Age
	≥ 12 years to < 18 years	≥ 18 years
Route	PO	PO
Dose	500-650 mg	960-1,000 mg
Max. single dose	650 mg	1,000 mg
Dosing interval	N/A	N/A
Max. # of doses	1	1

Consider ibuprofen		Consider ketorolac	
	Age		Age
	≥ 12 years		≥ 12 years
Route	PO	Route	IM/IV
Dose	400 mg	Dose	10-15 mg
Max. single dose	400 mg	Max. single dose	15 mg
Dosing interval	N/A	Dosing interval	N/A
Max. # of doses	1	Max. # of doses	1

Clinical Considerations

Whenever possible, consider co-administration of acetaminophen and ibuprofen.

Suspected renal colic patients should routinely be considered for NSAIDs, either ibuprofen or ketorolac.

IV administration of ketorolac applies only to PCPs authorized for PCP Autonomous IV.